

On the Classification of Finite Semigroups and RA -loops with the Hyperbolic Property

S. O. Juriaans^a A. C. Souza Filho^b

Instituto de Matemática e Estatística, Universidade de São Paulo, Caixa Postal 66281, São Paulo, CEP 05315-970 - Brazil

email addresses: ^aostanley@ime.usp.br ^bcalixto@ime.usp.br

Abstract

We classify the finite semigroups S , for which all $\mathbb{Z}$ -orders Γ of $\mathbb{Q}S$, the unit group $\mathcal{U}(\Gamma)$ is hyperbolic. We also classify the RA -loops L for which the unit loop of its integral loop ring does not contain any free abelian subgroup of rank two.

1 Introduction

Initially we classify finite dimensional algebras $\mathcal{A}$ over $\mathbb{Q}$ such that if $\Gamma \subset \mathcal{A}$ is a $\mathbb{Z}$ -order then $\mathcal{U}(\Gamma)$ is hyperbolic. If $\mathcal{A}$ is such an algebra, we say that $\mathcal{A}$ has the hyperbolic property.

In [7], are classified the semigroups Σ with $\mathcal{U}(\mathbb{Z}\Sigma)$ finite. Therefore, for this class of semigroups the algebra $\mathbb{Q}\Sigma$ has the hyperbolic property.

First we classify the semisimple algebras $\mathbb{Q}S$ which are nilpotent free. If $\mathbb{Q}S$ has nilpotent elements there are two possibilities: either S contains nilpotent elements, or not. In the latter case S is a disjoint union of groups of certain types. In the former, S is a union of groups and a subsemigroup of order five. We also give the structure of S when $\mathbb{Q}S$ is non-semisimple and has the hyperbolic property. For the proofs of the results of section two see [5].

In the last section we classify the RA -loops L , such that, $\mathbb{Z}^2 \not\curvearrowright \mathcal{U}(\mathbb{Z}L)$.

2 Semigroup Algebras

We will consider $\mathcal{A}$ a unitary finitely generated $\mathbb{Q}$ -algebra and denote by $\mathcal{S}(\mathcal{A})$, respectively $J(\mathcal{A})$, the semisimple subalgebra, respectively the Jacobson radical, of $\mathcal{A}$ and by $E(\mathcal{A}) = \{E_1, \dots, E_N\}$, $N \in \mathbb{Z}^+$, the set of the central primitive

idempotents of the semisimple algebra $\mathcal{S}(\mathcal{A})$. A classical result of Wedderburn-Mal'cev states that

$$\mathcal{A} \cong \mathcal{S}(\mathcal{A}) \oplus J(\mathcal{A}),$$

as a vector space. As a result, we have that $\mathcal{A}$ is a artinian algebra and thus its radical is a nilpotent ideal. We denote $T_2(\mathbb{Q}) := \begin{pmatrix} \mathbb{Q} & \mathbb{Q} \\ 0 & \mathbb{Q} \end{pmatrix}$ the 2×2 upper triangular matrices over $\mathbb{Q}$, with the usual matrix multiplication.

Definition 2.1 *Let $\mathcal{A}$ be a finite dimensional algebra over $\mathbb{Q}$ and Γ be a $\mathbb{Z}$ -order of $\mathcal{A}$. If*

$$\mathbb{Z}^2 \not\rightarrow \mathcal{U}(\Gamma),$$

we say $\mathcal{A}$ has the hyperbolic property.

Theorem 2.2 *Let $\mathcal{A}$ be a finite dimensional $\mathbb{Q}$ -algebra. If $\mathcal{A}_i$ is a simple epimorphic image of $\mathcal{A}$, denote by F_i a maximal subfield of $\mathcal{A}_i$ and $\Gamma_i \subset \mathcal{A}_i$ a $\mathbb{Z}$ -order. The following conditions hold:*

1. *The algebra $\mathcal{A}$ has the hyperbolic property, it is semisimple and it has no nilpotent element if, and only if,*

$$\mathcal{A} = \oplus \mathcal{A}_i,$$

whereof $\mathcal{A}_i$ is a division ring and there exists at most one index i_0 such that $\mathcal{U}(\Gamma_{i_0})$ is hyperbolic and infinite.

2. *The algebra $\mathcal{A}$ has the hyperbolic property and it is semisimple with nilpotent elements if, and only if,*

$$\mathcal{A} = (\oplus \mathcal{A}_i) \oplus M_2(\mathbb{Q}).$$

3. *The algebra $\mathcal{A}$ has the hyperbolic property and it is non semisimple with central radical if, and only if,*

$$\mathcal{A} = (\oplus \mathcal{A}_i) \oplus J.$$

4. *The algebra $\mathcal{A}$ has the hyperbolic property and it is non semisimple with non central radical if, and only if,*

$$\mathcal{A} = (\oplus \mathcal{A}_i) \oplus T_2(\mathbb{Q}).$$

For each item above, F_i is an imaginary quadratic field and $\mathcal{A}_i$ is either an imaginary quadratic field or a totally definite quaternion algebra. Furthermore, every simple epimorphic image of $\mathcal{A}$ in the direct sum is an ideal of $\mathcal{A}$.

In what follows, S denotes a finite semigroup, $\mathbb{Q}S$ denotes a unitary semigroup algebra over $\mathbb{Q}$, $\mathcal{M}^0(G; n, n; P)$ denotes the Rees semigroup with structural group G and P denotes an $n \times n$ sandwich matrix.

Theorem 2.3 *The algebra $\mathbb{Q}S$ has no nilpotent element and it has the hyperbolic property if, and only if, S is an inverse semi-group and it admits a principal series, whose principal factors are isomorphic to groups G and at most a unique K , listed below:*

1. G is an abelian group of exponent dividing 4 or 6;
2. G is a hamiltonian 2-group;
3. $K \in \{C_5, C_8, C_{12}\}$.

Theorem 2.4 *Let $\mathbb{Q}S$ be an algebra with nilpotent elements. The algebra $\mathbb{Q}S$ is semisimple and it has the hyperbolic property if, and only if, S admits a principal series whose principal factors are isomorphic to groups G and a unique semigroup K , listed below:*

1. G is an abelian group of exponent dividing 4 or 6.
2. G is a hamiltonian 2-group.
3. K is a group of the set $\{S_3, D_4, Q_{12}, C_4 \rtimes C_4\}$.
4. K is one of the Rees semigroups:

$$\mathcal{M}^0(\{1\}; 2, 2; I_d) = M \quad \text{or} \quad \mathcal{M}^0(\{1\}; 2, 2; \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}) = M_{12},$$

which is an ideal of S .

In particular, S is the disjoint union of the groups G and the semigroup K .

Theorem 2.5 *The algebra $\mathbb{Q}S$ is non semisimple and it has the hyperbolic property if, and only if, there exists a unique nilpotent element $j_0 \in S$, such that, the subsemigroup $\mathfrak{I} = \{\theta, j_0\}$ is an ideal of S , and $S \setminus \mathfrak{I}$ admits a principal series whose principal factors are isomorphic to abelian groups of exponent dividing 4 or 6, or a hamiltonian 2-group. In particular $S/\mathfrak{I}$ is the disjoint union of its maximal subgroups such that if $e_1 \in G_1$, and $e_N \in G_N$ are the respective group identity element, then $e_1 j_0 = j_0 e_N = j_0$. Writing*

$$\begin{aligned} e_1 &= \sum E_{1_i} + E_1 + \lambda j_0, \lambda \in \mathbb{Q} \\ e_N &= \sum E_{N_i} + E_N + \mu j_0, \mu \in \mathbb{Q} \end{aligned}$$

then only one of the following holds:

1.

$$e_1 e_N = 0 \Leftrightarrow e_N e_1 = 0 \text{ and } \lambda + \mu = 0;$$

$$T_2 \cong \{e_1, e_N, j_0, \theta\} \text{ is such that } \mathbb{Q}T_2 \cong T_2(\mathbb{Q}).$$

2.

$$\text{If } e_N e_1 \neq 0 \text{ then } e_1 e_N = e_N e_1 =: e_3 \text{ and } \lambda + \mu = 0;$$

$$T'_2 = \{e_1, e_2, e_3, j_0, \theta\} \text{ is a subsemigroup of } S \text{ and } \mathbb{Q}T'_2 \cong \mathbb{Q} \oplus \mathbb{Q} \oplus T_2(\mathbb{Q}).$$

3.

$$e_N e_1 = 0 \Leftrightarrow e_1 e_N = j_0 \Leftrightarrow \lambda + \mu = 1;$$

$$\hat{T}_2 = \{e_1, e_N, j_0, \theta\}, \text{ and } \mathbb{Q}\hat{T}_2 \cong T_2(\mathbb{Q}).$$

The semigroups T_2, T'_2 and $\hat{T}_2$ are non isomorphic.

3 The hiperbolicity of the RA -loop loop units

In this section we classify the RA -loops L such that $\mathbb{Z}^2 \not\rightarrow \mathcal{U}(\mathbb{Z}L)$, the loop of units of $\mathbb{Z}L$. A loop L is a nonempty set, with a closed binary operation $\cdot$, such that the equation $a \cdot b = c$ has a unique $b \in L$ when $a, c \in L$ are known, and a unique solution $a \in L$ when $b, c \in L$ are known, and with a two-side identity element 1. Denoting by $[x, y, z] = (xy)z - x(yz)$, recall that a ring A is alternative if $[x, x, y] = [y, x, x] = 0$, for every $x, y \in A$. An RA -loop is a loop whose loop ring RL over some commutative, associative and unitary ring R of characteristic not equal to 2 is alternative, but not associative. The basic reference is [10].

For a theoretical group property $\mathcal{P}$, a group G is virtually $\mathcal{P}$ if it has a subgroup of finite index, H say, with property $\mathcal{P}$.

Theorem 3.1 ([12], Theorem 3.3.6) *Let L be a RA -loop. $\mathcal{U}(\mathbb{Z}L)$ has the hyperbolic property if, and only if, L is a finite loop or a loop whose center is virtually cyclic, the torsion subloop $T(L)$ of L is such that, if $T(L)$ is a group, then it is an abelian group of exponent dividing 4 or 6 or a hamiltonian 2-group whose subgroups are all normal in L and if $T(L)$ is a loop then it is a hamiltonian Moufang 2-loop whose subloops are all normal in L . In this conditions we also have that $\mathcal{U}_1(\mathbb{Z}L) = L$.*

Acknowledgements

This work is part of the second authors Ph.D thesis, [12]. He would like to thank his thesis supervisor Prof. Dr. Stanley Orlando Juriaans for his guidance during this work.

References

- [1] L. G. X. de Barros, S. O. Juriaans, *Units in Alternative Integral Loop Rings*, Result. Math., 31 (1997), 266-281.
- [2] A. Borel, H. Chandra, *Arithmetic Subgroups of Algebraic Groups*, Annals of Mathematics, 75(3), 1962.
- [3] A. Doms, E. Jespers, *Generators for a subgroup of finite index in the unit group of an integral semigroup ring*, J. Group Theory 7(2004), 543-553.
- [4] M. Gromov, *Hyperbolic Groups*, in *Essays in Group Theory*, M. S. R. I. publ. 8, Springer, 1987, 75-263.
- [5] E. Iwaki, S. O. Juriaans, A. C. Souza Filho, *Hyperbolicity of Semigroup Algebras*, www.arxiv.org.
- [6] E. Jespers, *Free Normal Complements and the Unit Group of Integral Group Rings*, Proceedings of the American Mathematical Society, vol 122, number 1, 1994.
- [7] E. Jespers, D. Wang, *Units of Integral Semigroup Rings*, Journal of Algebra, vol 181, pages 395-413, 1996.
- [8] S. O. Juriaans, I. B. S. Passi, D. Prasad, *Hyperbolic Unit Groups*, Proceedings of the American Mathematical Society, vol 133(2), 2005, pages 415-423.
- [9] A. H. Clifford, G. B. Preston, *The Algebraic Theory of Semigroups*, American Mathematical Society, Mathematical Surveys number 7, Rhode Island, 1961.
- [10] E. G. Goodaire, E. Jespers, F. C. Polcino Milies, *Alternative Loop Rings*, Elsevier, Oxford, 1996.
- [11] J. Okniński, *Semigroup Algebras*, Pure and Applied Mathematics, Dekker, USA, 1991.
- [12] A.C. Souza Filho, *Sobre uma Classificação dos Anéis de Inteiros, dos Semigrupos Finitos e dos RA-Loops com a Propriedade Hiperbólica (On a Classification of the Integral Rings, Finite Semigroups and RA-Loops with the Hyperbolic Property)*, PhD. Thesis, IME-USP, São Paulo, 2006, 108 pages.